

PECTATE LYASE from Aspergillus sp. (Lot 111201d)

E-PCLYAN 02/17

(EC 4.2.2.2) (1->4)-alpha-D-galacturonan lyase

CAZy Family: PLI CAS: 9015-75-2

PROPERTIES

I. ELECTROPHORETIC PURITY:

- Single major band on SDS-gel electrophoresis (MW = ~ 45,000)

2. SPECIFIC ACTIVITY:

180 U/mg protein (on polygalacturonic acid) at pH 8.0 and 40°C

One Unit of pectate lyase activity is defined as the amount of enzyme required to release one µmole of galacturonic acid per minute from polygalacturonic acid (2.5 mg/mL) in Tris.HCl buffer (50 mM), pH 8.0 at 40°C.

3. SPECIFICITY:

Eliminative cleavage of (1,4)- α -D-galacturonan to give oligosaccharides with 4-deoxy- α -D-galact-4-enuronosyl groups at their non-reducing ends.

4. RELATIVE RATES OF HYDROLYSIS OF SUBSTRATES:

Substrate	%
Polygalacturonic acid (pectate lyase)	100 (Tris-Bispropane, pH 8.0)
Polygalacturonic acid (pectate lyase)	47.5 (CAPS buffer, pH 10.8)
Polygalacturonic acid (pectate lyase)	19.5 (Tris.HCl buffer, pH 8.0)
Polygalacturonic acid (endo-Polygalacturonanase)	< 0.0015
Galactazyme Tablets (endo-Galactanase)	< 0.0015
Arabinazyme Tablets (endo-Arabinanase)	< 0.0003

Action on polysaccharides was determined at a final substrate concentration 2.5 mg/mL in the relevant buffer. Activity was monitored at 235 nm in a recording spectrophotometer. *endo-*Polygalacturonanase, *endo-*arabinanase and *endo-*galactanase were assayed at pH 4.5 and 40°C.

5. PHYSICOCHEMICAL PROPERTIES:

pH Optima:	8.0
pH Stability:	6.5-11.0
Temperature Optima:	55°C
Temperature Stability:	< 55°C

6. STORAGE CONDITIONS:

The enzyme is supplied as a solution containing 50% glycerol and 0.02% (w/v) sodium azide and should be stored below -10°C. For assay, this enzyme should be diluted in Tris.HCl buffer (100 mM), pH 8.0. **Swirl to mix the enzyme immediately prior to use.**

The enzyme is also supplied as an ammonium sulphate suspension (E-PCLYAN2).

The enzyme is used for the identification of Pectin in foodstuffs, feed and fruit juices (K-PECID)