

INTERNAL STANDARD

OCHRATOXIN A

WHY

CHOOSE

LIBIOS?

Our LIB'UP® allows you to guarantee the quality of your analyses and accurately measure the toxins contained in your samples thanks to the high quality and purity of our standards.

If you use our internal standards, you avoid yield losses during your extractions, and you greatly reduce the matrix effect thanks to ion suppression.







Ready-to-use internal standard, certified with uncertainty for mycotoxins analysis by LC-MS/MS.



Uniformly enriched in C13, at 98% isotopic purity for an accurate and repeatable calibration.



Entire certificates in accordance with ISO 31, 34,35 and Eurachem / CITAC guides.



Tested and approved by external laboratories and French public research centers.



Various packaging: 0,5 mL - 1,2 mL - 5 mL - 10 mL and Custom mix.



Standards are designed, elaborated and produced by our laboratory in France.

OCHRATOXIN A DETERMINATION


The method increasingly used for the Ochratoxin A analysis is now the LC-MS/MS coupling. The analysis of this toxic exogenous compound requires the use of calibration reference solutions and / or the addition of internal standards as tracers of the molecule. The C13 labeled internal standard, Ochratoxin A, mimics as much as possible the physico-chemical behavior of the molecule to measure: identical structure and giving a specific and differentiated signal by the mass.

The mass spectrometry therefore allows the differentiation between the isotopologues and, by adding the known quantity of internal standard, the analyte content can be calculated. In other words, losses of the analyte during the purification various steps and / or extraction are completely compensated by similar losses of the isotopologue.


Ochratoxin A analysis example*:

*Test realised in our laboratory, use of our standard LIB'UP® U-[13C20]-Ochratoxin A.


MS spectrum of Ochratoxin A and standard LIB'UP® U-[13C20]-Ochratoxin A



Calibration curve obtained by using the standard LIB'UP® U-[13C20]-Ochratoxin A



High resolution scan of the LIB'UP® U-[13C20]-Ochratoxin A for determination of isotopic pattern



Product	Code/Packaging	Concentration	Solvent
[13C20]-Ochratoxin A	OTA13C20-10-0.5 (0.5 mL) OTA13C20-10-1.2 (1.2 mL) OTA13C20-10-5 (5 mL) OTA13C20-10-10 (10 mL)	10 μg/mL	Methanol



Phone: +33 (0)4 74 13 03 02

Mail: info@libios.fr Website: www.libios.fr

FLYER_IS_FUMONISIN_B3_EN_2022